

# POST-VACCINATION MANAGEMENT OF HIPRA'S ATTENUATED COCCIDIOSIS VACCINES

**EVALON**<sup>®</sup>  
With HIPRAMUNE<sup>®</sup> T

**EVANT**<sup>®</sup>  
With HIPRAMUNE<sup>®</sup> T

**EVANOVO**<sup>®</sup>

**Target: replication of attenuated vaccinal strains to build immunity against coccidiosis**

## 1 Environmental conditions




**23-28°C**  
(optimal conditions)



**30-35% Litter humidity**  
**55-60% Environmental humidity**

Tips for the evaluation of litter humidity



Video demo  
for humidity  
evaluation


## 2 Litter management

**These management strategies are only suggested if it is necessary to increase the litter humidity to 30 - 35% in the first 14 days.**


• Increased density of birds 

1-14 days	30-60 birds/m <sup>2</sup> (optimal) ≥ 20 birds/m <sup>2</sup> (minimum)
From 15 days	15-30 birds/m <sup>2</sup>

• Slightly increase water pipeline pressure  → 

• Irrigate lightly with a watering can / backpack once per day next to the pipelines 

• If the birds are moved to another area before 14 days, move **30%** of the litter to the new area  → 

• **Caged birds during first 3 weeks after vaccination: Paperboard or Cardboard** (220-250 gr./m<sup>2</sup>) a minimum of 21-24 days (50 - 100% of surface area) 

## 3 Warnings



Do not include the following substances at least during the cycling of the vaccine (first 3 weeks):

- Anticoccidial drugs (chemicals or ionophores)
- Amprolium, toltrazuril, sulfonamides or any coccidiocidal drug
- Tetracyclines and spiramycin
- Phytogenic compounds in high concentrations (>150 ppm)

## 4 Follow-up & sampling

**Vaccination day: administration procedure evaluation**

**7 days of age:** fresh faeces

**14 days of age:** fresh faeces/litter sample

**21 days of age:** fresh faeces/litter sample

**28 days of age:** litter sample

**35 days of age:** litter sample

Fresh faeces = pool of 20-30 fresh faeces

Litter sample = pool of 10 samples (25 grams per sample)

